



The MCD12 Products

- Primary inputs are MCD43A4/2 products
- MCD12Q1 and MCD12C1
 - Global, annual land cover at 500 m and 0.05°
 - 6 Types: IGBP, UMD, LAI/fPAR, NPP/BGC, PFT, LCCS
- MCD12Q2
 - Global, annual land surface phenology at 500 m
 - Start and end of season, integrated greenness



MCD12 Collection 6

- MCD12Q1/C1 and MCD12Q2 both use:
 - Daily Collection 6 NBARs
 - Spline smoothing of band time series
 - Snow screening and filling




MCD12Q1 Improvements

- Hierarchical classification approach
- RandomForest instead of C4.5
- Improved training data (STEP)
- Improved feature set
- Hidden Markov model (Abercrombie and Friedl, 2016 IEEE TGARS)



STEP Changes


47% more STEP sites
31% of sites modified or removed




Improved Algorithm and Classification Inputs

	V5	V6
Feature Set	 Monthly NBAR (7 bands), EVI, and LST. Annual metrics (min, max, mean) for each feature. 	 Annual metrics for growing season: from NBARs (7 bands) and vegetation indices (EVI2, NDII1, NDII2, NDWI, NDSI)
Classifier	C4.5 Decision Tree	RandomForest Classifier
Ancillary Data	 MODIS V5 land-water mask V5 urban layer Prior probability layers based on Collection 4 data with agriculture intensity data 	 MODIS V6 land-water mask at 250-m Updated V6 urban layer Prior probability layers based on: New agriculture intensity data Climate-based classifications MODIS Vegetation Continuous Fields product

New LCCS Layer








MCD12Q2

- What's new in C6?
 - Relative thresholds of splined NBAR-EVI2 rather than inversions of piecewise logistic functions
 - New SDS: peak, midpoint of GUP/GDOWN
 - Improved QA: QA value for each phenometric; weighted sum of missing data in vicinity of phenometric and spline goodness of fit











Phenometric-specific QA






QA score is weighted sum of fraction missing in month centered on phenometric and spline R^2 (80% and 20%, resp.) Continuous scores collapsed to 1-4 qualitative score


Extracting Phenology Data From Webcam Images





Harvard Forest Phenocam, MA











Kamuela Phenocam Waimea, HI


















50% Greenup Assessment with Melaas et al. Landsat method 2001-2014



Moving Forward

- Code migration to Goddard, will be processed on the PEATE like most other products
- Ongoing assessment and refinement
- Working with XYZ for VIIRS continuity



Questions?